



HOME & GARDEN

Potato or Tomato Psyllids

no. 5.540

by W.S. Cranshaw ¹

Quick Facts...

The potato/tomato psyllid secretes a toxic saliva during feeding that can severely damage potatoes and tomatoes.

Psyllids do not overwinter in Colorado. Outbreaks occur from flights of psyllids that migrate from southern states and Mexico.

Check potatoes and tomatoes each year for psyllid problems. If psyllids are detected, promptly treat affected plants with an insecticide.

Life Cycle

Potato/tomato psyllids pass through three life stages: egg, nymph (immature stage) and adult. The adult psyllid is about the size of a typical aphid and is a member of the insect family known as “jumping plant lice.” Adult psyllids are rarely found in gardens unless collected with a sweep net or knocked onto a cloth placed around the base of the plants. If seen, adult psyllids are striped with alternating dark and light bands (see Figure 1).

Eggs are small, 1/32 inch long. They are orange-yellow and supported by small stalks. They are much smaller than the stalked, white egg produced by lacewings, which also are common in gardens. Psyllid eggs are frequently deposited along leaf margins but may occur on either leaf surface. Eggs hatch in six to 10 days.

Newly hatched nymphs are yellowish but become progressively greener as they develop, undergoing four molts. When almost mature, nymphs are nearly the same color as leaves. Nymphs are flat, elliptical and scale-like.

Nymphs are most numerous on the undersides of leaves but can occur on shaded upper leaf surfaces. They are inactive and seldom can be seen to move about. While feeding, psyllid nymphs excrete small, waxy beads of “psyllid sugar,” which resembles granulated sugar. This material may cover leaves during heavy psyllid infestations. The nymph stage usually lasts from 14 to 22 days. Newly emerged adults remain green for a day or so before turning darker.

Potato/tomato psyllids do not overwinter in Colorado but survive on certain plants growing in southern Texas, New Mexico and Arizona. Problems with psyllids originate each season from winged, migrating forms of the insect. Psyllid outbreaks are irregular, depending on weather conditions.

Psyllids usually are found first on early potatoes or pepper transplants. Throughout the season, adult psyllids move to new plants, becoming most numerous late in the season on tomatoes. The number of psyllid generations produced during a year is thought to vary from four to seven. However, there is much overlap of the generations after the original infestations become established.




Figure 1: Tomato psyllid adult; tomato psyllid eggs on a leaf section; tomato psyllid nymph.



Figure 2: Psyllid nymphs feeding on potato leaf.




Figure 3: Psyllid feeding damage to potato plant.

Insect Injury

Adults and nymphs feed by sucking plant juices. Feeding by nymphs is especially serious because it brings about an abnormal condition known as “psyllid yellows,” a result of toxic saliva injected by the insect. The symptoms on potato and tomato plants are generally similar. Usually the first abnormal condition is a slight discoloration (yellowing or purpling) along the midribs and the edges of the top leaves. The basal portions of these leaves tend to curl upward.

As the condition progresses, the entire plant top changes to yellowish-green or purple-red, and foliar growth is checked. The leaves remain small and narrow and tend to stand upright, giving the top of the plant a feathery appearance.

When the attack comes early in the development of the tomato plant, effects from psyllid feeding may be so severe that little or no fruit is set. Late attack on tomato plants is inclined to cause production of an abnormal number of fruits that never attain a desirable size or quality.

If the attack on potato plants occurs before tuber set, a likely result is the formation of numerous tubers on each stolon. An attack after tubers are partially developed usually results in greatly retarded growth and irregularly shaped potatoes. Potatoes from infested plants may sprout prematurely, even underground before harvest.

Psyllids also occur on other plants in the potato family, such as eggplant and pepper. Damage to these crops is insignificant.

Control

Because these insects are so small, damage to tomatoes or potatoes frequently occurs before the problem is detected. It is important to be able to identify potato/tomato psyllids so developing problems can be detected and treated in time. One of the most important means of identification is the psyllid sugar that is excreted by the insect and collects on leaves. Psyllid problems do not occur every season. In some areas of the state, Cooperative Extension pest alerts provide warnings of psyllid outbreaks.

Homeowners not able to properly identify psyllids may wish to routinely treat susceptible plants. Protectant treatments may be needed from when plants are 6 inches tall until midsummer. Well-established plants with abundant foliage usually can tolerate late season infestations with little yield loss.

Among insecticides available to homeowners, Diazinon has proven most effective for psyllid control when used at rates for other potato/tomato insects. Thorough application is required for control. Repeat applications at 10- to 14-day intervals may be necessary if psyllids rapidly recolonize the plants. Preharvest interval requirements for Diazinon are 35 days on potatoes and one day on tomatoes.

Insecticidal soaps and diluted dishwashing detergents (1 to 2 percent concentration) also have been demonstrated to control psyllids. Apply soaps and detergents so that the psyllids are covered with the spray mixture. Some plants may be injured by soap or detergent sprays, so always test sprays first on a small area of the plant.

Some tomato varieties appear to be partially resistant to potato/tomato psyllids. Increased hairiness of the leaves is reported to make plants less favored by psyllids.